

**Announcements:**

Office hour Monday: 9:30 to 11 (or 10:55)  
Discussion Monday *is* for participation credit  
Weekly quiz will be available from today at  
2pm until Monday at 12:50.

**Homework** (due Monday):

Chapter 7: #47, 85



Chapter 7

**Probability  
continued**

**Review:**

**RELATIONSHIPS BETWEEN EVENTS**

- Defined for events in the *same random circumstance only*:
  - **Complement** of an event A is all simple events *not* part of A.
  - **Mutually exclusive = disjoint events** have no overlapping simple events
  - **Complements are disjoint events** (no overlapping simple events)
  - **Disjoint events aren't complements** *unless* they cover all possibilities.

**Review: Independent and Dependent Events**

- Two events are **independent** of each other if knowing that one will occur (or has occurred) *does not change* the probability that the other occurs.
- Two events are **dependent** if knowing that one will occur (or has occurred) *changes* the probability that the other occurs.

The definitions can apply *either ...*  
to events *within the same random circumstance* or  
to events *from two separate random circumstances*.

### EXAMPLE OF INDEPENDENT EVENTS

- Events in the *same random circumstance*:  
Daily 3 lottery on the *same* draw  
**A** = first two digits are 11  
**B** = last digit is 9  
 Knowing it starts with 11, P(B) is *still* 1/10.
- Events in *different random circumstances*:  
Daily 3 lottery on *different* draws  
**A** = today's winning number is 345  
**B** = tomorrow's winning number is 345  
 Knowing today's # was 345, P(B) is *still* 1/1000

### Conditional Probabilities

The **conditional probability** of the event **B**, **given that the event A occurs**, is the long-run relative frequency with which event B occurs when circumstances are such that A also occurs; written as  $P(B|A)$ .

$P(B)$  = unconditional probability event B occurs.

$P(B|A)$  = "probability of B given A"  
 = conditional probability event B occurs *given that we know A has occurred or will occur.*

*Illustrate with Venn diagram on board.*

### EXAMPLE OF CONDITIONAL PROBABILITY

TABLE 2.3 ■ Nighttime Lighting in Infancy and Eyesight

Slept with:	No Myopia	Myopia	High Myopia	Total
Darkness	155 (90%)	15 (9%)	2 (1%)	172
Nightlight	153 (66%)	72 (31%)	7 (3%)	232
Full Light	34 (45%)	36 (48%)	5 (7%)	75
Total	342 (71%)	123 (26%)	14 (3%)	479

*Random circumstance*: Observe one randomly selected child

**A** = child slept in darkness as infant

$$P(A) = 172/479 = .36$$

**B** = child did not develop myopia

$$P(B) = 342/479 = .71$$

$$P(B|A) = P(\text{no myopia} | \text{slept in dark}) = 155/172 = .90 \neq P(B)$$

*Illustrate with diagram on board.*

### NOTES ABOUT CONDITIONAL PROBABILITY

- $P(B|A)$  generally does *not* equal  $P(B)$ .
- When A and B are *independent* events,  $P(B|A) = P(B)$ , otherwise they are not equal.
- In Chapter 6, we were actually testing if two types of events were *independent*.
- Conditional* probabilities are similar to *row* and *column* proportions (percents) in contingency tables. (See myopia example on previous page.)

### HOW TO DETERMINE IF TWO EVENTS ARE INDEPENDENT

1. Physical assumption  
*Example:* Lottery draws on different days don't affect each other.
2. See if  $P(B|A) = P(B)$ . If so, A and B are independent events, otherwise they are not.  
*Example:* Suppose data showed that smokers and non-smokers are equally likely to get the flu.  
 $P(\text{flu} | \text{smoker}) = P(\text{flu})$  so they are independent.
3. Events A and B are independent if and only if  $P(A \text{ and } B) = P(A)P(B)$ , sometimes can check to see if this is true.

Copyright ©2004 Brooks/Cole, a division of Thomson Learning, Inc., updated by Jessica Uts, October 2009

## 7.4 Basic Rules for Finding Probabilities

### Probability an Event Does Not Occur

**Rule 1 (for “not the event”):**  $P(A^C) = 1 - P(A)$

**Example 7.13** *Probability a Stranger Does Not Share Your Birthday*

$P(\text{next stranger you meet will share your birthday}) = 1/365.$

$P(\text{next stranger you meet will not share your birthday}) = 1 - 1/365 = 364/365 = 0.9973.$

Copyright ©2004 Brooks/Cole, a division of Thomson Learning, Inc., updated by Jessica Uts, October 2009

### Probability That Either of Two Events Happen

**Rule 2 (addition rule for “either/or”):**

**Rule 2a (general):**

$$P(A \text{ or } B) = P(A) + P(B) - P(A \text{ and } B)$$

**Rule 2b (for mutually exclusive events):**

If A and B are *mutually exclusive* events,  
 $P(A \text{ or } B) = P(A) + P(B)$

*Illustrate with Venn diagram on board.*

Copyright ©2004 Brooks/Cole, a division of Thomson Learning, Inc., updated by Jessica Uts, October 2009

### Example 7.14 *Roommate Compatibility*

Brett is off to college. There are 1000 male students. Brett hopes his roommate will not like to party and not snore.

		Snores?		Total
		Yes	No	
Likes to Party?	Yes	150	100	250
	No	200	550	750
		350	650	1000

A = likes to party  $P(A) = 250/1000 = 0.25$

$P(A \text{ and } B) = 150/1000 = .15$

B = snores  $P(B) = 350/1000 = 0.35$

Probability Brett will be assigned a roommate who either likes to party or snores, or both is:  **$P(A \text{ or } B)$**

$$= P(A) + P(B) - P(A \text{ and } B) = 0.25 + 0.35 - 0.15 = 0.45$$

So the probability his roommate is acceptable is  $1 - 0.45 = 0.55$

## Probability That Two or More Events Occur Together

### Rule 3 (multiplication rule for “and”):

#### Rule 3a (general):

$$P(\mathbf{A} \text{ and } \mathbf{B}) = P(\mathbf{A})P(\mathbf{B}|\mathbf{A})$$

#### Rule 3b (for independent events):

If  $\mathbf{A}$  and  $\mathbf{B}$  are *independent* events,

$$P(\mathbf{A} \text{ and } \mathbf{B}) = P(\mathbf{A})P(\mathbf{B})$$

#### Extension of Rule 3b (> 2 independent events):

For several independent events,

$$P(\mathbf{A}_1 \text{ and } \mathbf{A}_2 \text{ and } \dots \text{ and } \mathbf{A}_n) = P(\mathbf{A}_1)P(\mathbf{A}_2)\dots P(\mathbf{A}_n)$$

Copyright ©2004 Brooks/Cole, a division of Thomson Learning, Inc., updated by Jessica Uts, October 2009

## Example with independent events (Rule 3b)

The probability of a birth being a boy is .512.  
Suppose a woman has 2 kids.

$\mathbf{A}$  = first child is a boy

$\mathbf{B}$  = second child is a boy

We assume these are independent events.

$$P(\mathbf{A} \text{ and } \mathbf{B}) = P(\mathbf{A})P(\mathbf{B}) = (.512)(.512) = .2621$$

$$\text{Probability of 2 girls is } (.488)(.488) = .2381$$

From **Rule 2b**, probability of same sex = .5002

From **Rule 1**, probability of different sex = .4998

## Example with dependent events (Rule 3a)

Randomly select a student who takes Stat 7 with me.

$\mathbf{A}$  = student comes to class regularly;  $P(\mathbf{A}) = 0.7$

$\mathbf{A}^C$  = student doesn't come regularly;  $P(\mathbf{A}^C) = 0.3$

$\mathbf{B}$  = student gets at least a B in the course

$$P(\mathbf{B}|\mathbf{A}) = 0.8 \quad P(\mathbf{B}|\mathbf{A}^C) = 0.4$$

What is the probability that the student comes to class regularly *and* gets at least a B?

$$P(\mathbf{A} \text{ and } \mathbf{B}) = P(\mathbf{A})P(\mathbf{B}|\mathbf{A}) = (.7)(.8) = .56$$

## Extension of Rule 3b:

### Example 7.17: Sharing Your Birth Month

Assume all 12 birth months are equally likely.

What is the probability that the next four *unrelated* strangers you meet *all share* your birth month?

**Event A** = 1<sup>st</sup> stranger shares your birth month  $P(\mathbf{A}) = 1/12$

**Event B** = 2<sup>nd</sup> stranger shares your birth month  $P(\mathbf{B}) = 1/12$

And so on, presumably all independent...

$$P(\text{four strangers share your birth month}) \\ = (1/12)(1/12)(1/12)(1/12) = (1/12)^4 = .000048$$

Copyright ©2004 Brooks/Cole, a division of Thomson Learning, Inc., updated by Jessica Uts, October 2009

## Determining a Conditional Probability

### Rule 4 (conditional probability):

$$P(B|A) = \frac{P(A \text{ and } B)}{P(A)}$$

$$P(A|B) = \frac{P(A \text{ and } B)}{P(B)}$$

Copyright ©2004 Brooks/Cole, a division of Thomson Learning, Inc., updated by Jessica Uts, October 2009

## EXAMPLE OF CONDITIONAL PROBABILITY - Revisited

TABLE 2.3 ■ Nighttime Lighting in Infancy and Eyesight

Slept with:	No Myopia	Myopia	High Myopia	Total
Darkness	155 (90%)	15 (9%)	2 (1%)	172
Nightlight	153 (66%)	72 (31%)	7 (3%)	232
Full Light	34 (45%)	36 (48%)	5 (7%)	75
Total	342 (71%)	123 (26%)	14 (3%)	479

A = child slept in darkness as infant

$$P(A) = 172/479 = .36$$

B = child did not develop myopia

$$P(B) = 342/479 = .71$$

$$P(B|A) = P(\text{no myopia} \mid \text{slept in dark}) = 155/172 = .90$$

But this is 
$$\frac{P(A \text{ and } B)}{P(A)} = \frac{155/479}{172/479} = \frac{155}{172}$$

Copyright ©2004 Brooks/Cole, a division of Thomson Learning, Inc., updated by Jessica Uts, October 2009

## Mutually exclusive versus Independent

Students sometimes confuse the definitions of **independent** and **mutually exclusive** events.

- When two events are *mutually exclusive* and one happens, it *turns the probability of the other one to 0*.
- When two events are *independent* and one happens, it *leaves the probability of the other one alone*.

Copyright ©2004 Brooks/Cole, a division of Thomson Learning, Inc., updated by Jessica Uts, October 2009

## In Summary (see page 249) ...

When Events Are:	$P(A \text{ or } B)$ is:	$P(A \text{ and } B)$ is:	$P(A B)$ is:
Mutually Exclusive	$P(A) + P(B)$	0	0
Independent	$P(A) + P(B) - P(A)P(B)$	$P(A)P(B)$	$P(A)$
Any	$P(A) + P(B) - P(A \text{ and } B)$	$P(A)P(B A)$	$\frac{P(A \text{ and } B)}{P(B)}$

Copyright ©2004 Brooks/Cole, a division of Thomson Learning, Inc., updated by Jessica Uts, October 2009

## 7.5 Strategies for Finding Complicated Probabilities

**Example 7.20** *Winning the Daily 3 Lottery*

Event A = winning number is 956. **What is  $P(A)$ ?**

**Method 1:** With physical assumption that all 1000 possibilities are equally likely,  $P(A) = 1/1000$ .

**Method 2:** Define three events,

$B_1 = 1^{\text{st}}$  digit is 9,  $B_2 = 2^{\text{nd}}$  digit is 5,  $B_3 = 3^{\text{rd}}$  digit is 6

Event A occurs if and only if all 3 of these events occur.

Note:  $P(B_1) = P(B_2) = P(B_3) = 1/10$ . Since these events are all *independent*, we have  $P(A) = (1/10)^3 = 1/1000$ .

\* *Can be more than one way to find a probability.*

## Steps for Finding Probabilities

**Step 1:** List each separate random circumstance involved in the problem.

**Step 2:** List the possible outcomes for each random circumstance.

**Step 3:** Assign whatever probabilities you can with the knowledge you have.

**Step 4:** Specify the event for which you want to determine the probability.

**Step 5:** Determine which of the probabilities from Step 3 and which probability rules can be combined to find the probability of interest.

Copyright ©2004 Brooks/Cole, a division of Thomson Learning, Inc., updated by Jessica Uts, October 2009

## Some Hints for Finding Probabilities

- **$P(A \text{ and } B)$ :** Sometimes you can define the event in physical terms and know the probability or find it from a two-way table.

Example: I could classify the class into male, female and also year in school. Then, for example, probability that a randomly selected student in the class is Male *and* sophomore is the proportion of the class in that cell of the table. Don't need separate  $P(A)$  and  $P(B)$ .

- Check if probability of the **complement** is **easier** to find, then subtract it from 1 (applying Rule 1).

Example: Probability of at least 1 boy in family of 3 kids =  
 $1 - \text{Probability of all girls} = 1 - (.488)^3 = 1 - .116 = .884$

Copyright ©2004 Brooks/Cole, a division of Thomson Learning, Inc., updated by Jessica Uts, October 2009

## Finding Conditional Probability in Opposite Direction

**Know  $P(B|A)$  but want  $P(A|B)$ :** Use Rule 3a to find  $P(B) = P(A \text{ and } B) + P(A^c \text{ and } B)$ , then use Rule 4.

$$P(A|B) = \frac{P(A \text{ and } B)}{P(B|A)P(A) + P(B|A^c)P(A^c)}$$

Let's look at some tools that are much easier than using the formula!

Copyright ©2004 Brooks/Cole, a division of Thomson Learning, Inc., updated by Jessica Uts, October 2009

## Solve these probability questions

### Example 1: Medical testing for a rare disease

D = person has the disease, suppose:

$$P(D) = 1/1000 = .001, P(D^c) = .999$$

T = test for the disease is positive, suppose:

$$P(T | D) = .95, \text{ so } P(T^c | D) = .05$$

$$P(T | D^c) = .05, \text{ so } P(T^c | D^c) = .95$$

So the test is **95% accurate** whether person has the disease or not

**FIND  $P(D | T)$**

= Probability of disease, *given* the test is positive

### Example 2: Probability of getting a B or better

Return to example of Statistics 7 grades

**A** = student comes to class regularly;  $P(A) = 0.7$

**B** = student gets at least a B in the course

$$P(B|A) = 0.8 \quad P(B|A^c) = 0.4$$

**Question:**

What is the overall probability of getting at least a B?

## Two-Way Table: "Hypothetical Hundred Thousand"

Table of hypothetical 100,000 people who get tested

1/1000 of them have disease = 100 people

Of those, 95% = 95 people test positive

99,900 don't have the disease

95% of those, or 94,905 people, test negative

So,  $P(\text{Disease} | \text{Test positive}) = 95/5090 = .019$

	Test pos	Test neg	
Disease	95	5	100
No disease	4995	94,905	99,900
	5090	94,910	100,000

Copyright ©2004 Brooks/Cole, a division of Thomson Learning, Inc., updated by Jessica Uts, October 2009

## Tree Diagrams

**Step 1:** Determine first random circumstance in sequence, and create first set of branches for possible outcomes. Create one branch for each outcome, write probability on branch.

**Step 2:** Determine next random circumstance and append branches for possible outcomes to each branch in step 1.

Write associated *conditional probabilities* on branches.


**Step 3:** Continue this process for as many steps as necessary.

**Step 4:** To determine the probability of following any particular sequence of branches, multiply the probabilities on those branches. This is an application of Rule 3a.

**Step 5:** To determine the probability of any collection of sequences of branches, add the individual probabilities for those sequences, as found in step 4. This is an application of Rule 2b.

Copyright ©2004 Brooks/Cole, a division of Thomson Learning, Inc., updated by Jessica Uts, October 2009

## Disease probability



$$P(D \text{ and positive test}) = .00095.$$

$$P(\text{test is positive}) = .00095 + .04995 = .0509.$$

$$P(D | \text{positive test}) = .00095 / .0509 = .019$$

Copyright ©2004 Brooks/Cole, a division of Thomson Learning, Inc., updated by Jessica Uits, October 2009


## Example 2: Return to example of Stat 7 grades

A = student comes to class regularly;  $P(A) = 0.7$

B = student gets at least a B in the course

$$P(B|A) = 0.8 \quad P(B|A^c) = 0.4$$

Question: What is the overall probability of getting at least a B?



So, probability of B or better =  $.56 + .12 = .68$  overall

More examples on board:

- Suppose there is no relationship between two variables, e.g. listening to Mozart and increased IQ. Suppose 3 independent experiments are done, each using the 0.05 criterion for statistical significance.

What is the probability that *at least one* finds statistical significance just by chance?

- Suppose you drive on a certain freeway daily. Speed limit is 65. You drive over 65 all the time, but over 75 about 30% of the time.

$$P(\text{ticket} | \text{over } 75) = 1/50 = .02$$

$$P(\text{ticket} | 65 \text{ to } 75) = 1/200 = .005.$$

What is the probability you get a ticket on a randomly selected day?