

# Universal Point Sets for Planar Graph Drawings with Circular Arcs



Patrizio Angelini, **David Eppstein**, Fabrizio Frati, Michael Kaufmann, Sylvain Lazard, Tamara Mchedlidze, Monique Teillaud, and Alexander Wolff

25th Canadian Conference on Computational Geometry  
Waterloo, Ontario, August 2013

# Fáry's theorem

Graphs that can be drawn with non-crossing curved edges can also be drawn with non-crossing straight edges

[Wagner 1936; Fáry 1948; Stein 1951]




...but not necessarily with the same vertex positions!

The set of points in  $\mathbb{R}^2$  is **universal** for straight drawings: it can be used to form the vertex set of any planar graph

# Smaller universal sets than the whole plane?

Every set of  $n$  points is universal for topological drawings  
(edges drawn as arbitrary curves) of  $n$ -vertex graphs




Simply deform the plane to  
move the vertices where you  
want them, moving the edges  
along with them

# Universal grids for straight line drawings

$O(n) \times O(n)$  square grids are universal  
[de Fraysseix et al. 1988; Schnyder 1990]

Some graphs require  $\Omega(n^2)$  area when drawn in grids



# Big gap for universal sets for straight line drawings


Best upper bound on universal point sets for straight-line drawing:

$$n^2/4 - O(n)$$

Based on *permutation patterns* [Bannister et al. 2013]

This 15-element permutation  
contains all 6-element  
213-avoiding permutations

Exponential stretching produces  
an 18-point universal set for  
9-vertex straight line drawings



Best lower bound:  $1.098n - o(n)$  [Chrobak and Karloff 1989]




## Two paths to perfection

**Perfect universal set:** exactly  $n$  points

Don't exist for straight drawings,  $n \geq 15$  [Cardinal et al. 2012]  
so have to relax either “straight” or “planar”.

Every  $n$ -point set in general position is universal for

- ▶ paths (connect in coordinate order)
- ▶ trees
- ▶ outerplanar graphs [Gritzmann et al. 1991]



What about drawing all planar graphs but relaxing straightness?

# Arc diagrams


Vertices placed on a line; edges drawn on one or more semicircles

Initially used for drawing nonplanar graphs with few crossings

[Saaty 1964; Nicholson 1968]


Later named and popularized in information visualization

[Wattenberg 2002]



# Monotone topological 2-page book embeddings

Every planar graph has a planar arc diagram with each edge drawn as a two-semicircle “S” curve [Giordano et al. 2007; Bekos et al. 2013]




- ▶ Add edges to make the graph maximal
- ▶ Find canonical order (each vertex above earlier ones, neighbors form contiguous path on upper boundary)
- ▶ Add each vertex to the right of its penultimate neighbor

(Useful property:  $\leq n - 1$  inflections between consecutive vertices)

# Perfect universal sets from monotone embeddings

Every  $n$  points on a line are universal for drawings in which edges are smooth curves formed from two circular arcs


Every set of  $n$  points is universal for polyline drawings with two bends per edge (mimic semicircles with steep zigzags)



Every smooth convex curve contains  $n$  points that are universal for polyline drawings with **one** bend per edge [Everett et al. 2010]

# Drawings with no bends and no inflections

What if we require each edge to be a single circular arc?



Lombardi drawing of a  
46-vertex non-Hamiltonian  
graph with cyclic edge  
connectivity five


[Grinberg 1968; Eppstein 2013]

Arc diagrams don't always exist and are NP-complete to find

Much recent interest in *Lombardi drawings* (evenly spaced edges at each vertex) [Duncan et al. 2012; Eppstein 2013] and *smooth orthogonal layouts* (axis-aligned arcs) [Bekos et al. 2013]

## Our result


For every  $n$ , there exists a perfect universal point set for drawings with circular-arc edges



Construction:

Choose  $n$  points on the parabola  $y = -x^2$   
at  $x$ -coordinates  $2^n, 2^{2n}, 2^{3n}, \dots, 2^{n^2}$

# How to draw a graph on this universal set




- ▶ Draw monotone topological book embedding
- ▶ Number vertices and inflection points from left to right, rounding vertex numbers up to multiples of  $n$
- ▶ Map point  $i$  to point on parabola with  $x = 2^i$
- ▶ Draw each edge as an arc through its three points

# Why is the resulting drawing planar?

Key properties, proved with some algebra:

Arc through any three points on parabola crosses it once from below to above  $\Rightarrow$  edges pass above/below vertices correctly



For six points  $x_0 \leq x_1 < x_2 < x_3 < x_4 \leq x_5$ , spaced exponentially, arcs  $x_0x_3x_4$  and  $x_1x_2x_5$  are disjoint  $\Rightarrow$  edges do not cross

# Conclusions

Perfect universal sets for circular-arc drawings

Purely a theoretical result—drawings are not usable

- ▶ Vertex placement requires exponential area
- ▶ Edges have very small angular resolution

In contrast, arc diagrams (with one arc per edge) are very usable and practical but can only handle a subset of planar graphs

Maybe some way of combining the advantages of both?

## References, I

- Michael J. Bannister, Zhanpeng Cheng, William E. Devanny, and David Eppstein. Superpatterns and universal point sets. In *Graph Drawing*, 2013. To appear.
- Michael A. Bekos, Michael Kaufmann, Stephen G. Kobourov, and Antonios Symvonis. Smooth orthogonal layouts. In *Graph Drawing 2012*, volume 7704 of *LNCS*, pages 150–161. Springer, 2013.
- Jean Cardinal, Michael Hoffmann, and Vincent Kusters. On Universal Point Sets for Planar Graphs. Electronic preprint arxiv:1209.3594, 2012.
- M. Chrobak and H. Karloff. A lower bound on the size of universal sets for planar graphs. *SIGACT News*, 20:83–86, 1989.
- Hubert de Fraysseix, János Pach, and Richard Pollack. Small sets supporting Fary embeddings of planar graphs. In *20th ACM Symp. Theory of Computing*, pages 426–433, 1988.
- Christian A. Duncan, David Eppstein, Michael T. Goodrich, Stephen G. Kobourov, and Martin Nöllenburg. Lombardi drawings of graphs. *J. Graph Algorithms and Applications*, 16(1):85–108, 2012.

## References, II

- David Eppstein. Planar Lombardi drawings for subcubic graphs. In *Graph Drawing 2012*, volume 7704 of *LNCS*, pages 126–137. Springer, 2013.
- Hazel Everett, Sylvain Lazard, Giuseppe Liotta, and Stephen Wismath. Universal Sets of  $n$  Points for One-Bend Drawings of Planar Graphs with  $n$  Vertices. *Discrete Comput. Geom.*, 43(2):272–288, 2010.
- István Fáry. On straight-line representation of planar graphs. *Acta Sci. Math. (Szeged)*, 11:229–233, 1948.
- Francesco Giordano, Giuseppe Liotta, Tamara Mchedlidze, and Antonios Symvonis. Computing upward topological book embeddings of upward planar digraphs. In *Algorithms and Computation (ISAAC 2007)*, volume 4835 of *LNCS*, pages 172–183. Springer, 2007.
- Ē. Ja. Grinberg. Plane homogeneous graphs of degree three without Hamiltonian circuits. In *Latvian Math. Yearbook 4*, pages 51–58. Izdat. “Zinatne”, Riga, 1968.
- P. Gritzmann, B. Mohar, János Pach, and Richard Pollack. Embedding a planar triangulation with vertices at specified positions. *Amer. Math. Monthly*, 98(2):165–166, 1991.

## References, III

- T. A. J. Nicholson. Permutation procedure for minimising the number of crossings in a network. *Proc. IEE*, 115:21–26, 1968.
- Thomas L. Saaty. The minimum number of intersections in complete graphs. *Proc. National Academy of Sciences*, 52:688–690, 1964.
- Walter Schnyder. Embedding planar graphs on the grid. In *1st ACM/SIAM Symp. Disc. Alg. (SODA)*, pages 138–148, 1990.
- S. K. Stein. Convex maps. *Proc. AMS*, 2(3):464–466, 1951.
- Klaus Wagner. Bemerkungen zum Vierfarbenproblem. *Jahresbericht der Deutschen Mathematiker-Vereinigung*, 46:26–32, 1936.
- M. Wattenberg. Arc diagrams: visualizing structure in strings. In *IEEE Symp. InfoVis*, pages 110–116, 2002.